

Christine Coillard  
Michel A. Leroux  
Karl F. Zabjek  
Charles Hilaire Rivard

## SpineCor – a non-rigid brace for the treatment of idiopathic scoliosis: post-treatment results

Received: 4 November 2000  
Revised: 27 May 2002  
Accepted: 7 June 2002  
Published online: 7 November 2002  
© Springer-Verlag 2002

This study was funded by the Research Centre, Sainte-Justine Hospital, Montréal, Québec.

C. Coillard · K.F. Zabjek  
C.H. Rivard (✉)  
Department of Surgery,  
Faculty of Medicine,  
University of Montréal, Québec, Canada  
e-mail: crivard@justine.umontreal.ca,  
Tel.: +1-514-3452344,  
Fax: +1-514-3454723

C. Coillard · M.A. Leroux · K.F. Zabjek  
C.H. Rivard  
Research Centre, Sainte-Justine Hospital,  
3175 Côte Ste-Catherine,  
Montréal, Québec, Canada H3T 1C5

**Abstract** The objective of this study was to assess the success of treatment during the follow-up of a group of 195 idiopathic scoliosis (IS) patients consecutively treated with the SpineCor system. A survival analysis was performed to estimate the cumulative probability of success during treatment, at follow-up and for the combined treatment and follow-up period. Success was defined as either a correction or stabilization of  $\pm 5^\circ$  or more, and failure as a worsening of more than  $5^\circ$ . The patient cohort was categorized before treatment into curves less than  $30^\circ$  (group 1), and curves greater than  $30^\circ$  (group 2). The survival analysis indicated a cumulative probability of success that increased during treatment with the patient wearing the brace (Year 1: 0.30, 0.39; Year 2: 0.62, 0.79; Year 3: 0.92, 0.89, for groups 1 and 2 respectively). During the post-treatment follow-up period, there was a stabilization (Year 1 post-treatment: 0.94, 0.89; Year 2 post-treatment: 0.85, 0.81), with an overall probability of success of 0.92 and 0.88 after

4 years of combined treatment and post-treatment follow-up. For the 29 patients who had a minimum follow-up of 2 years (initial Cobb angle:  $30^\circ \pm 9^\circ$ ), the trend during treatment was a decrease in spinal curvature at 3 months, with a mean difference of  $10^\circ$  (SD  $5^\circ$ ); at termination of treatment a mean difference of  $7^\circ$  (SD  $7^\circ$ ); and at the time of the 1- and 2-year follow-ups there was a difference of  $4^\circ$  (SD  $7^\circ$ ) and  $5^\circ$  (SD  $7^\circ$ ) respectively, with reference to the initial out of brace condition. At 2 years follow-up there was an overall correction of greater than  $5^\circ$  for 55% of the patients, 38% had a stabilisation and 7% had worsened by more than  $5^\circ$ . This initial cohort of patients demonstrated a general trend of initial decrease in spinal curvature in brace, followed by a correction and/or stabilisation at the end of treatment, which was maintained through 1, and 2 years' follow-up.

**Keywords** Idiopathic scoliosis · Orthopaedic treatment · Orthosis · Prognosis · Reducibility

### Introduction

The treatment of idiopathic scoliosis (IS) with a rigid brace used to be considered as either a therapeutic panacea, or excessive and disappointing, until it found its proper place in the range of treatment options [16, 18, 20]. Although there is evidence that rigid brace treatment is effective in altering the natural history of IS [11, 16], a significant

challenge still exists for clinicians to define an optimal treatment approach. It is recognised that factors such as the amplitude of the curvature [6, 10], the level of maturation [6, 10], and the maximum reducibility of the curvature [6, 14] are associated with post-treatment outcome. However, questions still remain regarding the optimal application of external forces [6, 14], and the consequence of rigid brace treatment on the integrity of the muscular system when a rigid brace is applied. The nature

of the challenge faced by clinicians arises from the multiple factors that contribute to curve progression [9], as well as the complex issue of optimally applying external forces to the spine to favour curve correction and stabilisation during periods of rapid growth and development [1, 8]. With this in consideration, as well as knowledge of the unique spinal region specific vertebral morphology [3, 12, 15] and mobility [19], a unique spinal curvature specific “corrective movement principle” [4, 5] was developed to correct and stabilise a spinal curvature, which is maintained and favoured by a non-rigid brace SpineCor [4, 5]. The SpineCor system also includes an anthropometrical and postural evaluation [21], which assists in the choice of classification, definition of the “corrective movement principle”, and the fitting of the SpineCor brace. The primary objective of this study is to evaluate the preliminary results of the first cohort of patients treated with the SpineCor system.

## Materials and methods

### Clinical study cohort

The therapeutic indication for treatment was based on a diagnosis of idiopathic scoliosis, where the patients demonstrated a progression of the Cobb angle of at least  $5^\circ$  confirmed by two X-rays at 6-month intervals. Other factors such as patient maturity, high growth potential, family history of surgery for severe idiopathic scoliosis, and a rib hump that is greater than  $7^\circ$  were also considered. Specific inclusion and exclusion criteria included the following:

### Inclusion criteria

- Idiopathic scoliosis diagnosis and radiographic confirmation of absence of significant pathological malformation of the spine
- Age 6–14 years old
- Initial Cobb angle equal to or above  $15^\circ$
- Initial Cobb angle equal to or less than  $50^\circ$
- Risser 0, 1, 2 or 3
- Scoliosis with suspected high risk of evolution (family history or other prognostic factor) or proven to be progressive (Cobb angle increase of  $5^\circ$  or more confirmed by two X-rays at 6-month intervals)

### Exclusion criteria

- Postural scoliosis: when a supine posteroanterior radiograph shows an almost complete reduction and there is a leg length discrepancy
- Patient inability to follow all the treatment instructions
- Presence of a congenital malformation of the spine or spina bifida aperta or spondylolisthesis
- Neuromuscular scoliosis


The following curve types were treated with the SpineCor brace: thoracic (mid and high,  $n=72$ ), thoracolumbar (with or without pelvic obliquity,  $n=58$ ), lumbar ( $n=22$ ) and double curves ( $n=43$ ). The presence of a hypo-kyphosis was not considered as a contraindication for treatment with the SpineCor System.

The initial pre-therapeutic radiograph, which was used as a reference, was systematically taken following the classic method, us-

ing a digital technique where the irradiation is half as much as that of a standard radiograph [17]. The initial evaluation included a posteroanterior and lateral X-ray without brace within a maximum of 1 month before the brace was fitted. The following X-ray controls were always administered with the SpineCor brace following the same schedule: the first control on the day of the fitting and at 6 weeks and 3 months, then every 5 months on average until wearing. The lateral X-rays were obtained once a year. At the end of the treatment, the controls are continued at a rate of once every 6 months to 2 years, depending on the age of the child. These evaluations were performed without the brace on the patient.

### Description of the bracing system and treatment protocol

The dynamic corrective SpineCor brace resembles a non-rigid harness, and was developed at Sainte-Justine Hospital between 1992 and 1993. It consists of a pelvic base, which is a belt that includes three pieces of soft thermodeformable plastic stabilised by two thigh bands and two crotch bands, a bolero made of cotton and four corrective elastic bands of variable size (0.20–1 m). It is important to note that there are a number of configurations possible for the placement of the elastic bands. The therapeutic principle is based on the definition of a specific corrective movement for each type of curve [4, 5], with an adjustment of the corrective bands to reproduce and favour this corrective movement (Fig. 1, Fig. 2). The patients were requested to wear the brace 20 out of 24 h. The brace is stopped near skeletal maturity, or after 2 years of regular menstruation.



**Fig. 1** The corrective movement principle for a patient with a right thoracic curve. This movement involves a rotation of the thorax in a counter-clockwise direction relative to the shoulders (A, B) clockwise rotation of the shoulders and a slight down tilt of the right shoulder (A). The right lateral shift of T1 in relation to S1 should also be reduced (C)



**Fig. 2** The SpineCor brace fitted on a patient with a right thoracic curve, to favour the corrective movement principle

#### Statistical analysis

Patient visits were defined in the following manner. “Initial state” defines the patient’s status prior to treatment, “treatment 3 months (3 M)” defines the patient’s status in brace at 3 months into treatment, “end of treatment (ET)” is the evaluation date without brace when the weaning commenced, “follow-up 1 year (F1), 2 years (F2), 3 years (F3), 4 years (F4)” is defined as the out of brace follow-up. More specifically, differences (Cobb angle, and percentage change) between the initial condition and each following condition (Difference = Initial – 3 M, Initial – ET, Initial – F1, Initial – F2) as well as the changes that occurred during treatment and follow-up (Difference = 3 M – ET, 3 M – F1, 3 M – F2, ET – F1, ET – F2, F1 – F2) were analysed. The difference between visits was used to identify a success, defined as either an improvement of more than 5° or stabilisation of  $\pm 5^\circ$ , or a failure, defined by an aggravation of the spinal curvature of more than 5°.

The results were analysed such that an appreciation could be obtained for the general trend in treatment for the patients that had a minimum of 2 years follow-up. An indication of the efficacy of the SpineCor brace was obtained by performing a survival analysis including all of the patients in the cohort. A repeated measures analysis of variance was performed on the patients who completed treatment, to define the overall treatment trend of patients with a minimum follow-up of 2 years. Since this type of analysis will only include patients with an available visit at each time interval (Initial state, 3 M, ET, F1 and F2), the results of 29 patients with a minimum of 2 years follow-up were presented. Therefore, the patients who completed treatment but had a follow-up of less than 2 years, and those who withdrew prematurely or progressed to surgery were not included in the repeated measures analysis of variance. These patients were included in the following survival analysis.

The survival analysis is similar to the approach used by Nachemson and Peterson [11]. The whole cohort of patients was divided according to the amplitude of the initial Cobb angle such that group 1 (G1) consisted of patients with a Cobb angle less than 30° and group 2 (G2) consisted of patients with a Cobb angle greater than 30°. The criterion for success was defined as a correction or stabilization of the Cobb angle, and for failure as an aggravation of the Cobb angle. With the initial visit as a reference point, survival curves were constructed for:

1. The patients who were still under treatment as well as the withdrawals (Survival Analysis A)
2. The patients who had completed treatment, which includes surgical patients (Survival Analysis B)

A third survival curve was constructed with the end of treatment status as a reference point, and the last available visit during post-treatment follow-up as reference points. This analysis included all patients who completed treatment, including surgical patients (Survival Analysis C).

## Results

For this group of consecutively treated idiopathic scoliosis patients, the average age at the commencement of treatment was 13 years (SD 1 year); there were 176 female and 19 male subjects. The initial major Cobb angle for the patients with a major curve of less than 30° was 23° (SD 5°,  $n=115$ ), and for patients with a major curve of greater than 30° the Cobb angle was 36° (SD 4°,  $n=80$ ). The initial cohort characteristics by curve amplitude and curve type as well as the minimum Cobb angle during treatment are presented in Table 1, and the Risser sign in Table 2.

At the last available visit, there were 109 patients still under treatment, with a mean treatment time of 1.5 years (SD 1 year), and 71 had terminated treatment, with a post-treatment follow-up time ranging from 0 to 4.5 years, including eight patients who underwent surgery. The patients who progressed to surgery had an initial mean Cobb angle of 34° (SD 5°), with a mean end of treatment Cobb angle of 45° (SD 6°) after 2 years (SD 1 year) of treatment. There were also 15 patients who withdrew from treatment after a mean of 1.2 years (SD 0.72 years), for whom non-compliance and re-location were the principal reasons.

#### Outcome for idiopathic scoliosis patients with a minimum 2-year follow-up after treatment with the SpineCor system

From the 71 patients who completed treatment, there were 29 who had a minimum follow-up time of 2 years (mean 29 months, SD 4 months), and 42 who did not have a minimum of 2 years follow-up. The initial Cobb angle for this sub-cohort of patients was 29° (SD 7°), and after 3 months of treatment the mean Cobb angle was 19° (SD 11°), corresponding to an overall mean decrease of 10° (SD 5°), representing a mean reducibility of 40% (SD 28%). At the end of treatment (time = 24 months; SD 9 months; Risser


**Table 1** Initial characteristics of the idiopathic scoliosis patient population (group 1: initial Cobb angle less than 30°, group 2: initial Cobb angle more than 30°)

	Initial Cobb angle (°)		Cobb angle minimum during treatment (°)		Percent reduction (%)	
	Mean	SD	Mean	SD	Mean	SD
<b>All patients (n=195)</b>	29	8	18	10	38	26
Thoracic (n=72)	30	8	20	10	35	24
Thoracolumbar (n=58)	25	8	13	8	50	26
Lumbar (n=22)	24	6	16	7	36	25
Double (n=43)	32	7	23	11	31	26
<b>Group 1 (n=115)</b>	23	5	13	8	45	28
Thoracic (n=37)	24	4	15	7	41	26
Thoracolumbar (n=44)	22	4	10	7	54	27
Lumbar (n=18)	22	4	15	7	36	27
Double (n=16)	24	5	15	9	42	32
<b>Group 2 (n=80)</b>	36	4	26	8	28	20
Thoracic (n=35)	36	4	26	8	28	21
Thoracolumbar (n=14)	36	4	23	6	35	19
Lumbar (n=4)	33	2	21	6	37	18
Double (n=27)	37	4	28	8	23	18

**Table 2** Risser sign for the idiopathic scoliosis patient population

	R0	R1	R2	R3	R4
Group 1	86	7	12	10	0
Group 2	46	12	7	13	2

**Fig. 3** Results of patients weaned off the SpineCor, with a minimum 2 year follow-up



3 or 4), the mean Cobb angle was 21° (SD 12°), at 1 year follow-up it was 25° (SD 11°) and at 2 years follow-up it was 24° (SD 11°).

At 2 years follow-up there was an overall correction in reference to the patient's initial state of greater than 5° for 16 patients [mean: 10°; range: 6° (16%) to 15° (83%)]. All of these patients had a reduction at 3 months of greater than 5°, representing a mean reducibility of 51% (SD 28%). At 2 years follow-up, there was a stabilisation for 11 patients [mean 2° (9%); range -3° (19%) to 5° (25%)]. Of these patients, six had an initial reduction of less than 5°, which was maintained through to follow-up, and five patients had an initial reduction of greater than 5°, which was lost by 2 years follow-up. There were two patients who worsened [mean: -8° (31%); range: -6° (-17%) to -10° (-33%)] at 2 years follow-up. Both patients had an initial reduction of greater than 5°, which was lost during treatment. The evolution during treatment for the improved, stabilised and aggravated patients are presented in Fig. 3.

#### General treatment trend

For the 29 patients who had a minimum of 2 years follow-up, a repeated measures analysis of variance was performed, comparing the initial state, 3 months in brace, end of treatment, 1 year and 2 years follow-up. Since this is a preliminary analysis of an initial cohort of 29 patients, a significance level of  $P < 0.01$  was chosen. There was a significant difference between the initial condition and 3 months ( $10^\circ \pm 5^\circ$ ), end of treatment ( $7^\circ \pm 7^\circ$ ) as well as follow-up at 1 and 2 years ( $4^\circ \pm 7^\circ$  and  $5^\circ \pm 7^\circ$ ), respectively. There was no difference between 3 months and end of treatment ( $-3^\circ \pm 8^\circ$ ), and 1-year follow-up ( $-6^\circ \pm 7^\circ$ ), but

**Table 3** Difference between each time interval during the course of treatment with the SpineCor system and during follow-up (3M 3 months, ET end of treatment, 1Y 1 year follow-up, 2Y 2 year follow-up)

	Initial – 3M, ET, 1Y, 2Y				3 M – ET, 1Y, 2Y			ET – 1Y, 2Y		1Y – 2Y
	3M	ET	1Y	2Y	ET	1Y	2Y	1Y	2Y	
All	10 (5)	7 (7)	4 (7)	5 (7)	-3 (8)	-6 (7)	-5 (7)	-3 (6)	-2 (5)	1 (4)
Improved	12 (4)	10 (6)	8 (5)	9 (3)	-2 (4)	-4 (5)	-2 (3)	-2 (7)	-1 (5)	2 (5)
Stable	7 (6)	5 (5)	0 (3)	1 (3)	-2 (9)	-6 (5)	-5 (7)	-4 (6)	-4 (6)	1 (3)
Aggravated	10 (3)	-7 (1)	-13 (1)	-13 (1)	-17 (1)	-23 (1)	-23 (1)	-6 (0)	-6 (0)	0

there was a difference between 3 months and the 2-year follow-up ( $-5^{\circ}\pm 7^{\circ}$ ). Compared with the end of treatment, there was no difference with 1 year follow-up ( $-3^{\circ}\pm 6^{\circ}$ ), or with 2 year follow-up ( $-2^{\circ}\pm 5^{\circ}$ ), nor was there any difference between the 1 year and 2 year follow-ups ( $1^{\circ}\pm 4^{\circ}$ ) (see Table 3 and Table 4).

**Survival analysis of patients treated with the SpineCor system**

*Survival analysis A (n=124): patients under treatment and withdrawals (initial visit in reference to the last available visit)*

The cumulative probability of success at 1 year, 2 years and 3 years in treatment is presented in Table 5 and Fig. 4. In general, the probability of obtaining a positive treatment effect increased as the duration of treatment increased, for both groups of patients. There was a significant difference between the patients of group 1 (patients with an initial Cobb angle less than 30°) and those of group 2 (those with an initial Cobb angle greater than 30°) as identified by the log rank test ( $P=0.03$ ).


**Table 4** Repeated measures analysis of variance for patients with a minimum follow-up of 2 years

Conditions	P-value
Initial vs 3 months in brace	0.0000*
Initial vs end of treatment	0.0003*
Initial vs 1 year follow-up	0.0002*
Initial vs 2 years follow-up	0.0006*
3 months vs end of treatment	0.0135
3 months vs 1 year follow-up	0.0181
3 months vs 2 years follow-up	0.0081*
End of treatment vs 1 year follow-up	0.999 <sup>a</sup>
End of treatment vs 2 years follow-up	0.999 <sup>a</sup>
1 year vs 2 years follow-up	0.999 <sup>a</sup>

\*  $P<0.01$   
<sup>a</sup> The comparison between end of treatment and 1 year and 2 year follow-up indicates that there is no worsening of the curve during the post-treatment follow-up period. This is in contrast with most rigid braces

**Table 5** Cumulative probability of success of patients under treatment, including withdrawals, with the SpineCor system

Time	Group 1	Group 2
1 year	0.30 (CI: 0.18–0.41)	0.39 (CI: 0.25–0.52)
2 years	0.62 (CI: 0.49–0.75)	0.79 (CI: 0.67–0.90)
3 years	0.92 (CI: 0.84–1.0)	0.89 (CI: 0.80–0.99)




**Fig. 4** Cumulative probability of failure. The survival function for 124 SpineCor patients under treatment (including withdrawals) indicates that the probability of failure decreases with time in both group 1 (patients with an initial Cobb angle of less than 30°) and group 2 (those with an initial Cobb angle of more than 30°). (1 represents 100% probability of failure. The probability of success can be calculated by: 1 – prob failure)

*Survival Analysis B (n=71): all patients who terminated treatment (global success at follow-up: initial visit in reference to last available visit)*

Since the average treatment time is 2 years, the cumulative probability of success was calculated at 3, 4 and 5 years after the fitting of the SpineCor system, with the last available visit out of brace for the weaned patients, which included the surgical patients. As presented in Table 6 and Fig.5, the cumulative probability of success for group 1

**Table 6** Cumulative probability of success of patients who have completed treatment in reference to the initial fitting of the SpineCor system

Time	Group 1	Group 2
3 years	0.97 (CI: 0.91–1.0)	0.95 (CI: 0.87–1.0)
4 years	0.92 (CI: 0.82–1.0)	0.88 (CI: 0.71–1.0)
5 years	0.69 (CI: 0.44–0.93)	0.89 (CI: 0.62–1.0)

**Fig. 5** Survival function for 71 SpineCor patients who completed treatment, in reference to their initial status. The probability of success ranges from 0.97 at 3 years to 0.69 at 5 years for group 1, and from 0.95 at 3 years to 0.89 at 5 years for group 2. Note: the results at 5 years represents a small number of patients as reflected by the size of the confidence interval


and group 2 patients at 3 years follow-up was 0.97 and 0.95 respectively, and at 4 years follow-up it was 0.92 and 0.88 respectively. At 5 years there was a considerable decrease; however, at this time the confidence interval is two times that at 3 and 4 years. This is most likely due to the limited number of patients available for the analysis during this time period. There was no significant difference between the patients of group 1 and group 2 as identified by the log rank test.

*Survival analysis C (n=71): all patients who terminated treatment (global success at follow-up: end of treatment in reference to last available visit)*

The cumulative probability of success was calculated at 1, 2, and 3 years follow-up in reference to the end of treatment status with the SpineCor system. As shown in Table 7 and Fig. 6, the cumulative probability of success decreased as the duration of post-treatment follow-up period

**Table 7** Cumulative probability of success of patients for the duration of time between the end of treatment and the last available follow-up visit

Time	Group 1	Group 2
1 year	0.94 (CI: 0.86–1.0)	0.89 (CI: 0.84–1.0)
2 years	0.85 (CI: 0.71–0.98)	0.81 (CI: 0.62–1.0)
3 years	0.68 (CI: 0.36–0.99)	–

**Fig. 6** Survival function for SpineCor patients who completed treatment in reference to their end of treatment status. The probability of success ranges from 0.94 at 1 year to 0.68 at 3 years for group 1, and from 0.89 at 1 year to 0.81 for group 2, at 1 and 2 years. Note: the results at 3 years represent a small number of patients, as demonstrated by the size of the confidence interval

increased. However, since there is considerable overlap between the confidence intervals at 1 and 2 years follow-up, a stabilisation can also be a possible interpretation. At 3 years there was a considerable decrease. However, this is most likely due to the limited number of patients available for the analysis during this time period. There was no significant difference between the patients of group 1 and group 2 as identified by the log rank test.

## Discussion

The primary objective of this study was to perform a preliminary evaluation of the long-term outcome results of the first patients who completed treatment with the SpineCor System. For the 29 patients who had a minimum follow-up of 2 years, there was an overall correction greater than 5° for 55% of the patients, 38% had a stabilisation and 7% had worsened by more than 5°. The trend during treatment was to have a decrease in spinal curva-

ture at 3 months with a mean difference of  $10^\circ$  (SD  $5^\circ$ ), at termination of treatment a mean difference of  $7^\circ$  (SD  $7^\circ$ ), and at a follow-up time of 1 and 2 years a difference of  $4^\circ$  (SD  $7^\circ$ ) and  $5^\circ$  (SD  $7^\circ$ ) respectively, with reference to the initial out of brace condition. The survival analysis indicated an increasing cumulative probability of success over time for the patients in treatment in brace, which increased from 0.30 (group 1) and 0.39 (group 2) at 1 year, to 0.92 (group 1) and 0.89 (group 2) at 3 years in treatment. For the weaned patients, the cumulative probability of success was 0.85 (group 1) and 0.81 (group 2) at 2 years after the brace stopped being worn, with a global probability of 0.92 (group 1) and 0.88 (group 2) at 4 years after the initial fitting of the brace.

These results are similar to those reported previously by other bracing systems [2, 6, 10]. However, the principal difference is noted to involve the amplitude of initial correction and the amplitude of the final correction obtained at follow-up. Rigid bracing systems have been reported to have a maximum in-brace reducibility that ranges from 50 to 62% [6, 14]. A reducibility of greater than 50% is associated with a maintained correction that can reach  $7.2^\circ$  at follow-up [6, 7, 10, 14], and a reducibility of 8–10% is associated with a failure of the brace [13]. In the present study, the mean reducibility for all patients was 38%, which is lower than that reported by other bracing systems [6, 7, 10, 14]. However, if the patients are categorised according to those who had an overall correction, stabilisation or worsening at 2 years follow-up, the results are very similar to other bracing systems [6, 7, 10, 14]. With the SpineCor system there was a significant correlation ( $r=-0.73$ ) between the reducibility at 3 months and the amplitude of spinal curvature at 2 years follow-up. For the patients who had a decrease in spinal curvature of greater than  $5^\circ$  at 2 years follow-up, the mean correction was  $10^\circ$  and the initial reducibility at 3 months was 51%. For the patients whose status did not change by more than or less than  $5^\circ$  at follow-up, representing a mean change of  $2^\circ$ , the initial reducibility at 3 months was 27%. For the patients who deteriorated by more than  $5^\circ$  at follow-up with a mean aggravation of  $10^\circ$ , there was an initial reducibility of 30% at 3 months. These results support the notion that the more the curve is reduced during the brace treatment, the better are the chances of correction and stabilisation [6, 7, 10, 14]. However, when deliberating using the initial reducibility as a global indicator, it is also important to bear in mind that in some cases this notion does not apply. For example, in this study, a patient with an initial low reducibility of 25% had a final correction of  $12^\circ$ , which is the converse of the result of another patient, who had a strong initial reducibility (50%) but had an eventual aggravation of the Cobb angle of  $8^\circ$ . These exceptions may be attributed to the presence of significant vertebral deformation as well as the difficulty encountered by all bracing systems in controlling the curve during periods of rapid growth. The difference in the am-

plitude of reducibility between SpineCor and other bracing systems may be related to the treatment principles used. Rigid braces rely on a more direct three-point pressure principle in contrast to the SpineCor system, which involves the “corrective movement principle”. The corrective movement acts indirectly on the spinal column, and allows some degree of controlled mobility and movement. This approach provides the opportunity to re-educate and maintain the neuromuscular control of spinal corrective movement through active bio-feedback.

At the end of treatment, the patients with a minimum of 2 years follow-up in this study demonstrated a mean out-of-brace Cobb angle that was  $5^\circ$  lower than the initial pre-treatment status. The mean out of brace Cobb angle at termination of treatment was similar to that found for other bracing systems, where a correction of  $1^\circ$ – $4^\circ$  has been noted [6, 10]. In the present study, 70% of the patients maintained their correction/stabilisation from the end of treatment up until 2 years follow-up. It is hypothesized by the authors that the controlled mobility and movement that is allowed by the SpineCor system contributes to maintaining neuromuscular system integrity as well as educating neuromuscular control patterns to function in a favourable manner. Once the brace is discontinued, the corrective movement is integrated by the neuromuscular system, and the change in spinal curvature is maintained through follow-up. This may also account for the patients who showed a small improvement during follow-up. However, in some instances there was deterioration in the spinal curvature between the end of treatment and follow-up. For example, for three patients there was a deterioration of  $6^\circ$ ,  $7^\circ$ , and  $9^\circ$  respectively. Two patients had a Risser 3 and one patient a Risser 4 at the termination of treatment, with a Cobb angle of  $28^\circ$ ,  $18^\circ$  and  $18^\circ$  respectively. During the 1st year of post-treatment follow-up, the respective Risser signs were 5, 4 and 5, with a loss in correction of  $3^\circ$ ,  $4^\circ$  and  $7^\circ$  respectively. During the 2nd year of follow-up, with a Risser sign at 5 for all three patients, the first patient lost an additional  $6^\circ$ , the second  $2^\circ$  and the third remained the same. The angular loss during the 1st year post-treatment may be related to the possibility that these three patients were not fully matured. However, for a patient who experienced an angular loss of  $6^\circ$  during the 2nd year, despite going from Risser 4 to 5, the loss in curvature may no longer be related to growth, but mostly to factors that include deformation of the vertebrae, condition of the discs, muscular imbalance and overall tonus. These factors may have been present early on in treatment for this patient, who at the initiation of treatment had a Cobb angle of  $38^\circ$ . The difficulty in maintaining the correction and stabilisation of curves close to  $40^\circ$  has also been encountered by other bracing systems [6, 7].

The definition of bracing as an intermediate treatment option has largely relied on the comparison of the natural history of the disease with progression of the curvature during bracing treatment. A non-treated curve with a

Risser sign between 0 and 1 and a Cobb angle between 20° and 29° had a 68% chance of progressing [9]. In a meta-analysis based on 20 studies, the weighted mean proportion of success was determined to be 0.93 for full time bracing [16]. A comparative study between electrical stimulation, bracing and natural history identified that bracing was associated with a success rate of 74% at 4 years, which was statistically superior to observation only (34%), and electrical stimulation (33%) [11]. The results of the present evaluation of the SpineCor System infers an equally important if not greater success rate as reported by other bracing systems. At 4 years, the global probability of success was 0.92 and 0.88 for Cobb angles less than 30° and greater than 30° respectively. At a follow-up of 2 years, there was an improvement for 55%, stabilisation for 38%, and aggravation for 7%. However, a limitation of the present study is that the results are based on patients treated consecutively with the SpineCor system for all

type of curves. A more direct comparison to a non-treated control group as well as a group treated with a rigid bracing system would provide a stronger basis for evaluating the efficacy of the SpineCor System, and is the focus of ongoing investigations.

## Conclusion

This initial cohort of idiopathic scoliosis patients who were treated with the SpineCor system reveals a positive treatment outcome at 2 years follow-up. This is reflected through a cumulative probability of success, which increases during treatment, and is maintained through 1 and 2 years follow-up. For the patients followed from the initiation of treatment through to 2 years follow-up, there was an overall correction/stabilisation for 93% of the patients.

## References

- Aubin CE, Labelle H, Ruszkowski A, Petit Y, Gignac D, Joncas J, Dansereau J (1999) Variability of strap tension in brace treatment for adolescent idiopathic scoliosis. *Spine* 24:349–354
- Carr WA, Moe JH, Winter RB, Lonstein JE (1980) Treatment of idiopathic scoliosis in the Milwaukee brace. Long-term results. *J Bone Joint Surg Am* 62:599–612
- Coillard C, Rivard CH (1996) Vertebral deformities and scoliosis. *Eur Spine J* 5:91–100
- Coillard C, Leroux MA, Badeaux J, Rivard CH (2000) SpineCor: a new therapeutic approach for idiopathic scoliosis. Presented at the Annual Eurospine Meeting, Anvers, 10–14 October
- Coillard C, Leroux MA, Badeaux J, Rivard CH (2000) SpineCor: a new therapeutic approach for idiopathic scoliosis. Presented at the Bi-Annual Meeting of the International Research Society of Spinal Deformities (IRSSD) Clermont-Ferrand, 26–30 May
- Emans JB, Kaelin A, Bancel P, Hall JE, Miller ME (1986) The Boston bracing system for idiopathic scoliosis. Follow-up results in 295 patients. *Spine* 11:792–801
- Katz DE, Richards S, Browne RH, Herring JA (1997) A comparison between the Boston brace and the Charleston Bending Brace in adolescent idiopathic scoliosis. *Spine* 22:1302–1312
- Labelle H, Dansereau J (1990) Orthotic treatment of pediatric spinal disorders and diseases. *Spine State of the Art Reviews* 4:239–251
- Lonstein JE, Carlson JM (1984) The prediction of curve progression in untreated idiopathic scoliosis during growth. *J Bone Joint Surg Am* 66:1061–1071
- Lonstein JE, Winter RB (1994) The Milwaukee brace for the treatment of adolescent idiopathic scoliosis. A review of one thousand and twenty patients. *J Bone Joint Surg Am* 76:1207–1221
- Nachemson AL, Peterson L (1995) Effectiveness of treatment with a brace in girls who have adolescent idiopathic scoliosis. *J Bone Joint Surg Am* 77:6
- Nguyen VH, Leroux M, Badeaux J, Zabjek K, Coillard C, Rivard CH (1998) Classification des scoliozes thoraco-lombaires gauches selon leur morphologie radiologique et leur géométrie posturale. *Ann Chir* 52:752–760
- Noonan K, Weinstein SL, Jacobson WC, Dolan LA (1996) Use of the Milwaukee brace for progressive idiopathic scoliosis. *J Bone Joint Surg Am* 78:557–567
- Olafsson Y, Saraste H, Soderlund V, Hoffsten M (1995) Boston brace in the treatment of idiopathic scoliosis. *J Pediatr Orthop* 15:524–527
- Rivard CH, Coillard C, Badeaux J, Leroux M, Zabjek KF, Simard G (2000) Idiopathic scoliosis. New classification, how and why. Example: Thoracolumbar curves. Fifth Annual Meeting of the Société internationale de recherche et d'étude du rachis (SIRER), Rome, 23–24 November
- Rowe DE, Bernstein SM, Riddick MF, Adler F, Emans JB, Gardner-Bonneau D (1997) A meta-analysis of the efficacy of non-operative treatments for idiopathic scoliosis. *J Bone Joint Surg Am* 79:664–674
- Roy M, Boutard A, Labelle H (1997) La radiologie numérique permet-elle vraiment de diminuer l'exposition aux radiations chez les adolescents avec scoliose idiopathique? Conference Proceedings of the Annual Meeting of the Communication Société de la scoliose du Québec
- Schuffelbarger HL, Keiser RP, King W (1983) Nonoperative treatment of idiopathic scoliosis: a 10-year study. *Orthop Trans* 7:11–13
- White AA, Panjabi MM (1990) *Clinical biomechanics of the spine*, 2nd edn. Lippincott, Philadelphia, pp 128–168
- Winter RB (1994) The pendulum has swung too far. *Orthop Clin North Am* 25:195–205
- Zabjek KF, Simard G, Leroux MA, Coillard C, Rivard CH (1999) Comparison of the reliability of two 3D acquisition systems used for the study of anthropometric and postural parameters. *Ann Chir* 53:751–760